

MEASURE NAME	MEASURE DESCRIPTION	eMEASUR	eMEASUR	NGF	QUALITY	NGS DOMS	MEASURE	HIGH Prio	DATA SUBMISSION METHOD	SPECIALTY MEASURE SET	PRIMARY MEASURE STEWARD
Annual Hepatitis C Virus (HCV) Screening for Patients who are Active Injection Drug Users	Percentage of patients, regardless of age, who are active injection drug users who were screened for HCV infection within the 12 month reporting period	None	None	None	387	ECC	Process	No	Registry	Internal Medicine, General Practice/Family Medicine	Physician Consortium for Performance Improvement
Anti-Depressant Medication Management	Percentage of patients 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression, and who remained on an antidepressant medication treatment. Two rates are reported: a. Percentage of patients who remained on an antidepressant medication for at least 84 days (12 weeks). b. Percentage of patients who remained on an antidepressant medication for at least 180 days (6 months).	CMS128v	None	105	9	ECC	Process	No	EHR	Internal Medicine, Mental/Behavioral Health, General Practice/Family Medicine	National Committee for Quality Assurance
Atrial Fibrillation and Atrial Flutter: Chronic Anticoagulation Therapy	Percentage of patients aged 18 years and older with a diagnosis of nonvalvular atrial fibrillation (AF) or atrial flutter whose assessment of the specified thromboembolic risk factors indicates one or more high-risk factors or more than one moderate risk factor, as determined by CHADS2 risk stratification, who are prescribed warfarin OR another oral anticoagulant drug that is FDA approved for the prevention of thromboembolism	None	None	1525	326	ECC	Process	No	Claims, Registry	Internal Medicine, Cardiology, General Practice/Family Medicine	American College of Cardiology
Breast Cancer Screening	Percentage of women 50-74 years of age who had a mammogram to screen for breast cancer.	CMS125v	None	2372	112	ECC	Process	No	Claims, EHR, CMS Web Interface, Registry	Internal Medicine, Obstetrics/Gynecology, Preventive Medicine, General Practice/Family Medicine	National Committee for Quality Assurance
Cervical Cancer Screening	Percentage of women 21-64 years of age who were screened for cervical cancer using either of the following criteria: * Women age 21-64 who had cervical cytology performed every 3 years * Women age 30-64 who had cervical cytology/human papillomavirus (HPV) co-testing performed every 5 years	CMS124v	None	32	309	ECC	Process	No	EHR	Obstetrics/Gynecology, General Practice/Family Medicine	National Committee for Quality Assurance
Colorectal Cancer Screening	Percentage of adults 50-75 years of age who had appropriate screening for colorectal cancer.	CMS130v	None	34	113	ECC	Process	No	Claims, EHR, CMS Web Interface, Registry	Internal Medicine, General Practice/Family Medicine	National Committee for Quality Assurance
Coronary Artery Disease (CAD): Beta-Blocker Therapy-Prior Myocardial Infarction (MI) or Left Ventricular Systolic Dysfunction (LVEF <40%)	Percentage of patients aged 18 years and older with a diagnosis of coronary artery disease seen within a 12 month period who also have a prior MI or a current or prior LVEF <40% who were prescribed beta-blocker therapy	CMS145v	None	70	7	ECC	Process	No	EHR, Registry	Cardiology, General Practice/Family Medicine	Physician Consortium for Performance Improvement
Diabetes: Eye Exam	Percentage of patients 18-75 years of age with diabetes who had a retinal or dilated eye exam by an eye care professional during the measurement period or a negative retinal exam (no evidence of retinopathy) in the 12 months prior to the measurement period	CMS131v	None	55	112	ECC	Process	No	Claims, CMS Web Interface, EHR, Registry	Internal Medicine, Ophthalmology, General Practice/Family Medicine	National Committee for Quality Assurance
Diabetes: Foot Exam	The percentage of patients 18-75 years of age with diabetes (type 1 and type 2) who received a foot exam (visual inspection and sensory exam with mono filament and a pulse exam) during the measurement year	CMS123v	None	56	163	ECC	Process	No	EHR	Internal Medicine, General Practice/Family Medicine	National Committee for Quality Assurance
Diabetes: Medical Attention for Nephropathy	The percentage of patients 18-75 years of age with diabetes who had a nephropathy screening test or evidence of nephropathy during the measurement period.	CMS134v	None	62	119	ECC	Process	No	EHR, Registry	General Practice/Family Medicine	National Committee for Quality Assurance
Documentation of Signed Opioid Treatment Agreement	All patients 18 and older prescribed opiates for longer than six weeks duration who signed an opioid treatment agreement at least once during Opioid Therapy documented in the medical record.	None	None	None	412	ECC	Process	No	Registry	Internal Medicine, Neurology, Physical Medicine, General Practice/Family Medicine	American Academy of Neurology
Evaluation or Interview for Risk of Opioid Misuse	All patients 18 and older prescribed opiates for longer than six weeks duration evaluated for risk of opioid misuse using a brief validated instrument (e.g. Opioid Risk Tool, SOAPP-R) or patient interview documented at least once during Opioid Therapy in the medical record	None	None	None	414	ECC	Process	No	Registry	Internal Medicine, Neurology, Physical Medicine, General Practice/Family Medicine	American Academy of Neurology
Heart Failure (HF): Angiotensin-Converting Enzyme (ACE) Inhibitor or Angiotensin Receptor Blocker (ARB) Therapy for Left Ventricular Systolic Dysfunction (LVSD)	Percentage of patients aged 18 years and older with a diagnosis of heart failure (HF) with a current or prior left ventricular ejection fraction (LVEF) <40% who were prescribed ACE inhibitor or ARB therapy either within a 12 month period when seen in the outpatient setting OR at each hospital discharge	CMS135v	2907	81	5	ECC	Process	No	EHR, Registry	Internal Medicine, Cardiology, Hospitalists, General Practice/Family Medicine	Physician Consortium for Performance Improvement
Heart Failure (HF): Beta-Blocker Therapy for Left Ventricular Systolic Dysfunction (LVSD)	Percentage of patients aged 18 years and older with a diagnosis of heart failure (HF) with a current or prior left ventricular ejection fraction (LVEF) <40% who were prescribed beta-blocker therapy either within a 12 month period when seen in the outpatient setting OR at each hospital discharge	CMS144v	2908	83	8	ECC	Process	No	EHR, Registry	Cardiology, Hospitalists, General Practice/Family Medicine	Physician Consortium for Performance Improvement
Hepatitis C: Screening for Hepatocellular Carcinoma (HCC) in Patients with Cirrhosis	Percentage of patients 18 years and older with a diagnosis of chronic hepatitis C cirrhosis who underwent imaging with either ultrasound, contrast enhanced CT or MRI for hepatocellular carcinoma (HCC) at least once within the 12 month reporting period	None	None	None	401	ECC	Process	No	Registry	Internal Medicine, Gastroenterology, General Practice/Family Medicine	American Gastroenterological Association
Immunizations for Adolescents	The percentage of adolescents 13 years of age who had the recommended immunizations by their 13th birthday	None	None	1407	394	CPH	Process	No	Registry	General Practice/Family Medicine, Pediatrics	National Committee for Quality Assurance
Ischemic Vascular Disease (IVD): Use of Aspirin or Another Antiplatelet	Percentage of patients 18 years of age and older who were diagnosed with acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous coronary interventions (PCI) in the 12 months prior to the measurement period, or who had an active diagnosis of ischemic vascular disease (IVD) during the measurement period, and who had documentation of use of aspirin or another antiplatelet during the measurement period.	CMS154v	None	68	204	ECC	Process	No	Claims, EHR, CMS Web Interface, Registry	Internal Medicine, Cardiology, General Practice/Family Medicine	National Committee for Quality Assurance
One-Time Screening for Hepatitis C Virus (HCV) for Patients at Risk	Percentage of patients aged 18 years and older with one or more of the following: a history of injection drug use, receipt of a blood transfusion prior to 1992, receiving maintenance hemodialysis, OR birthdate in the years 1945-1965 who received one-time screening for hepatitis C virus (HCV) infection	None	None	None	400	ECC	Process	No	Registry	Internal Medicine, General Practice/Family Medicine	Physician Consortium for Performance Improvement
Opioid Therapy Follow-up Evaluation	All patients 18 and older prescribed opiates for longer than six weeks duration who had a follow-up evaluation conducted at least every three months during Opioid Therapy documented in the medical record	None	None	None	408	ECC	Process	No	Registry	Internal Medicine, Neurology, Physical Medicine, General Practice/Family Medicine	American Academy of Neurology
Osteoporosis Management in Women Who Had a Fracture	The percentage of women age 50-85 who suffered a fracture and who either had a bone mineral density test or received a prescription for a drug to treat osteoporosis in the six months after the fracture	None	None	53	418	ECC	Process	No	Claims, Registry	Internal Medicine, Obstetrics/Gynecology, General Practice/Family Medicine	National Committee for Quality Assurance
Persistence of Beta-Blocker Treatment After a Heart Attack	The percentage of patients 18 years of age and older during the measurement year who were hospitalized and discharged from July 1 of the year prior to the measurement year to June 30 of the measurement year with a diagnosis of acute myocardial infarction (AMI) and who were prescribed persistent beta-blocker treatment for six months after discharge	None	None	71	442	ECC	Process	No	Registry	General Practice/Family Medicine	National Committee for Quality Assurance
Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan	Percentage of patients aged 18 years and older with a BMI documented during the current encounter or during the previous six months AND with a BMI outside of normal parameters, a follow-up plan is documented during the encounter or during the previous six months of the current encounter Normal Parameters: Age 18 years and older BMI >= 18.5 and < 25 kg/m2	CMS69v	None	421	128	CPH	Process	No	Claims, EHR, CMS Web Interface, Registry	Internal Medicine, Cardiology, Gastroenterology, General Surgery, Hospitalists, Neurology, Obstetrics/Gynecology, Orthopedic Surgery, Otolaryngology, Physical Medicine, Preventive Medicine, Rheumatology, Vascular Surgery, Mental/Behavioral Health, Plastic Surgery, General Practice/Family Medicine	Centers for Medicare & Medicaid Services
Preventive Care and Screening: Influenza Immunization	Percentage of patients aged 6 months and older seen for a visit between October 1 and March 31 who received an influenza immunization OR who reported previous receipt of an influenza immunization	CMS147v	None	41	110	CPH	Process	No	Claims, EHR, CMS Web Interface, Registry	Allergy/Immunology, Internal Medicine, Obstetrics/Gynecology, Preventive Medicine, General Practice/Family Medicine, Pediatrics	Physician Consortium for Performance Improvement
Preventive Care and Screening: Screening for Clinical Depression and Follow-Up Plan	Percentage of patients aged 12 years and older screened for depression on the date of the encounter using an age appropriate standardized depression screening tool AND if positive, a follow-up plan is documented on the date of the positive screen	CMS216	None	418	134	CPH	Process	No	Claims, EHR, CMS Web Interface, Registry	Internal Medicine, Mental/Behavioral Health, General Practice/Family Medicine, Pediatrics	Centers for Medicare & Medicaid Services
Preventive Care and Screening: Screening for High Blood Pressure and Follow-Up Documented	Percentage of patients aged 18 years and older seen during the reporting period who were screened for high blood pressure AND a recommended follow-up plan is documented based on the current blood pressure (BP) reading as indicated	CMS225v	None	None	312	CPH	Process	No	Claims, EHR, Registry	Allergy/Immunology, Internal Medicine, Cardiology, Dermatology, Emergency Medicine, Gastroenterology, General Surgery, General Oncology, Hospitalists, Neurology, Obstetrics/Gynecology, Ophthalmology, Orthopedic Surgery, Otolaryngology, Physical Medicine, Preventive Medicine, Rheumatology, Thoracic Surgery, Urology, Vascular Surgery, Mental/Behavioral Health, Plastic Surgery, General Practice/Family Medicine	Centers for Medicare & Medicaid Services
Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention	Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received cessation counseling intervention if identified as a tobacco user	CMS138v	None	28	226	CPH	Process	No	Claims, EHR, CMS Web Interface, Registry	Internal Medicine, Cardiology, Emergency Medicine, Gastroenterology, General Oncology, Hospitalists, Neurology, Obstetrics/Gynecology, Ophthalmology, Orthopedic Surgery, Otolaryngology, Physical Medicine, Preventive Medicine, Rheumatology, Thoracic Surgery, Urology, Vascular Surgery, Mental/Behavioral Health, Plastic Surgery, General Practice/Family Medicine	Physician Consortium for Performance Improvement
Preventive Care and Screening: Unhealthy Alcohol Use: Screening & Brief Counseling	Percentage of patients aged 18 years and older who were screened for unhealthy alcohol use using a systematic screening method at least once within the last 24 months AND who received brief counseling if identified as an unhealthy alcohol user	None	None	2152	431	CPH	Process	No	Registry	Internal Medicine, Mental/Behavioral Health, General Practice/Family Medicine	Physician Consortium for Performance Improvement

Statin Therapy for the Prevention and Treatment of Cardiovascular Disease	Percentage of the following patients-all considered at high risk of cardiovascular events-who were prescribed or were on statin therapy during the measurement period: Adults aged >= 21 years who were previously diagnosed with or currently have an active diagnosis of clinical atherosclerotic cardiovascular disease (ASCVD); OR adults >="190" (ldl-c)="" a="" active="" aged="" an="" cholesterol="" currently="" diagnosed="" diagnosis="" direct="" dl="" ever="" familial="" fasting="" had="" have="" hypercholesterolemia;="" level="" li="" lipoprotein="" low-density="" mg="" of="" or="" or<="" previously="" pure="" were="" who="" with="" years="">adults 40-75="" 70-189="" a="" aged="" diabetes="" diagnosis="" direct="" dl.<="" fasting="" ldl-c="" level="" li="" mg="" of="" or="" with="" years="">Noneadults>adults>	None	None	438	ECC	Process	No	CMS Web Interface,Registry	Internal Medicine,Cardiology,General Practice/Family Medicine	Centers for Medicare & Medicaid Services	
Tobacco Use and Help with Quitting Among Adolescents	The percentage of adolescents 12 to 20 years of age with a primary care visit during the measurement year for whom tobacco use status was documented and received help with quitting if identified as a tobacco user	None	None	None	402	CPH	Process	No	Registry	Allergy/Immunology,Internal Medicine,Cardiology,Dermatology,Emergency Medicine,Gastroenterology,General Surgery,General Oncology,Hospitalists,Neurology,Obstetrics/Gynecology,Ophthalmology,Orthopaedic Surgery,Otolaryngology,Physical Medicine,Preventive Medicine,Rheumatology,Thoracic Surgery,Urology,Vascular Surgery,Mental/Behavioral Health,Plastic Surgery,General Practice/Family Medicine,Pediatrics	National Committee for Quality Assurance
Tuberculosis (TB) Prevention for Psoriasis, Psoriatic Arthritis and Rheumatoid Arthritis Patients on a Biological Immune Response Modifier	Percentage of patients whose providers are ensuring active tuberculosis prevention either through yearly negative standard tuberculosis screening tests or are reviewing the patient's history to determine if they have had appropriate management for a recent or prior positive test	None	None	None	337	ECC	Process	No	Registry	Dermatology,Rheumatology,General Practice/Family Medicine	American Academy of Dermatology